

The sub-millimetre evolution of V4334 Sgr (Sakurai's Object)

A. Evans¹, T. R. Geballe², V. H. Tyne¹, D. Pollacco³, S. P. S. Eyres⁴,
B. Smalley¹

¹*Department of Physics, Keele University, Keele, Staffordshire, ST5 5BG*

²*Gemini Observatory, 670 N. A'ohoku Place, Hilo, HI 96720*

³*Department of Pure & Applied Physics, Queen's University of Belfast, Belfast, BT7 1NN*

⁴*Centre for Astrophysics, University of Central Lancashire, Preston, PR1 2HE*

Revised version: 22 July

ABSTRACT

We report the results of monitoring of V4334 Sgr (Sakurai's Object) at $450\,\mu\text{m}$ and $850\,\mu\text{m}$ with SCUBA on the James Clerk Maxwell Telescope. The flux density at both wavelengths has increased dramatically since 2001, and is consistent with continued cooling of the dust shell in which Sakurai's Object is still enshrouded, and which still dominates the near-infrared emission. Assuming that the dust shell is optically thin at sub-millimetre wavelengths and optically thick in the near-infrared, the sub-millimetre data imply a mass-loss rate during 2003 of $\sim 3.4 \pm 0.2 \times 10^{-5} \text{ M}_{\odot} \text{ yr}^{-1}$ for a gas-to-dust ratio of 75. This is consistent with the evidence from $1 - 5\,\mu\text{m}$ observations that the mass-loss is steadily increasing.

Key words: stars, individual: V4334 Sgr – stars, individual: Sakurai's Object – stars: evolution

1 INTRODUCTION

V4334 Sgr (Sakurai's Object) continues to be the subject of intense interest. It is a low-mass ($\lesssim 0.7 \text{ M}_{\odot}$) star that is retracing its evolution along the Hertzsprung-Russell diagram following a very late thermal pulse (Herwig 2001). The star is at the centre of a faint planetary nebula (PN) of diameter $40''$ (e.g. Pollacco 1999; Bond & Pollacco 2002) and, after an early incarnation as an F-type star, it underwent a transformation to a carbon star in 1997.

Little is known observationally about the evolution of the star since mid-1998, when it was completely obscured by an optically thick dust shell from which (as of April 2004) it has still not emerged; a dramatic illustration of the obscuration is given by Bond & Pollacco (2002). Since the major dust event of 1998 the $1 - 5\,\mu\text{m}$ spectrum is black-body-like and is simply that of the dust shell. Analysis of the post-1998 infrared (IR) observations has shown that the dust is carbon, primarily in amorphous form (Geballe et al. 2002; Tyne et al. 2002; Eyres et al., in preparation). The $1 - 5\,\mu\text{m}$ spectra during the period 1999–2001 were consistent with a mass-loss that increased from $\sim 1.8 \times 10^{-6} \text{ M}_{\odot} \text{ yr}^{-1}$ to $\sim 4.4 \times 10^{-6} \text{ M}_{\odot} \text{ yr}^{-1}$ (for the gas-to-dust ratio of 75 we adopt here; see §3 below), while the maximum grain radius in a grain size distribution $n(a) da \propto a^{-q} da$ ($q \simeq 3$) increased by a factor ~ 3 (Tyne et al. 2002).

We first observed V4334 Sgr at sub-millimetre wave-

lengths with the SCUBA instrument (Holland et al. 1999) on the 15-m James Clerk Maxwell Telescope (JCMT) in 2001 August, and reported marginal detections at both $450\,\mu\text{m}$ and $850\,\mu\text{m}$ (Evans et al. 2002, hereafter Paper I).

We report here the results of our continued monitoring of Sakurai's Object at these wavelengths.


2 OBSERVATIONS AND DATA REDUCTION

We have observed Sakurai's Object on the dates given in Table 1, in which we include our 2001 August observation from Paper I. On each occasion atmospheric extinction was regularly monitored by taking skydip measurements, and pointing sources in the observatory catalogue were used to check telescope pointing and focussing; Uranus and Mars were used as calibrators. The beamwidths at $450\,\mu\text{m}$ and $850\,\mu\text{m}$ are $7''$ and $14''$ respectively.

SCUBA was used in PHOTOMETRY mode, which provides simultaneous observations at $450\,\mu\text{m}$ and $850\,\mu\text{m}$; the observations were carried out in good (sub-millimetre) weather conditions to ensure that data at both $450\,\mu\text{m}$ and $850\,\mu\text{m}$ were obtained. However, as is well-known (Holland et al. 1999), radiation at the shorter wavelength is severely affected by atmospheric water vapour, and hence detection and calibration is always more problematic at $450\,\mu\text{m}$ even when weather conditions are favourable. The observations

Table 1. Sub-millimetre flux densities f_ν of V4334 Sgr. Errors are 1σ .

Date D/M/Y	MJD – 2450000	FCF (mJy mV ^{−1})		f_ν (450 μ m) (mJy)	Detection level (σ)	f_ν (850 μ m) (mJy)	Detection level (σ)	Comment
		450 μ m	850 μ m					
18/08/2001	2140	309 $\pm$ 72	219 $\pm$ 30	96.6 $\pm$ 37.5	3.2	10.6 $\pm$ 3.4	3.5	Paper I
09/01/2003	2649	378 $\pm$ 49	193 $\pm$ 5	55.2 $\pm$ 14.9	4.2	35.7 $\pm$ 2.4	17	
05/05/2003	2765	322 $\pm$ 3	202 $\pm$ 1	157.6 $\pm$ 25.8	4.1	52.1 $\pm$ 2.7	20	
15/06/2003	2806	440 $\pm$ 5	217 $\pm$ 2	213.2 $\pm$ 29.1	7.4	61.3 $\pm$ 2.9	28	
16/06/2003	2807	401 $\pm$ 2	213 $\pm$ 1	113 $\pm$ 41	2.8	58.3 $\pm$ 3.2	18	

**Figure 1.** The 450 μ m (left) and 850 μ m (right) light curves of Sakurai's Object.

consisted of 2 (or 4) sets of 50 integrations, to give total on-source integration time of 30 (or 60) minutes. First order sky subtraction was achieved by chopping 60'' in azimuth.

The data were reduced using the SURF package (Jenness & Lightfoot 2000). The frames were flat-fielded and corrected for atmospheric extinction, and the outer bolometers in the array were used for second order sky subtraction; 'spikes' in the data were clipped to remove any points further than 3σ from the mean. The flux conversion factors at 850 μ m and 450 μ m, together with the detection levels (the signal-to-noise ratios in the measured signal) for Sakurai's Object are listed in Table 1.

Sakurai's Object was strongly detected at both 450 μ m and 850 μ m, and the flux densities f_ν are listed in Table 1. As usual, a significant fraction of the uncertainty in the flux densities for Sakurai's Object listed in Table 1 arises from the uncertainty in the flux calibration, particularly at 450 μ m. In the Table the flux errors include both statistical errors and calibration uncertainties. The measurement at 450 μ m for 2003 June 16 is marginal and is not used in what follows.

3 DISCUSSION

The 450 μ m and 850 μ m light curves are shown in Fig. 1. We see a consistent and significant rise in the 850 μ m flux density between 2001 August and 2003 September; the behaviour of the 450 μ m data is consistent with the rise at 850 μ m, but of course the 450 μ m calibration is less certain. In Paper I we considered the possibility that the 450/840 μ m emission detected in 2001 August might have originated in the old PN but we concluded that the emission originated in material ejected in the 1995 event. The fact that the sub-

millimetre flux densities rise so dramatically is consistent with this conclusion.

We first note that the sub-millimetre data may imply a drop in the β -index of the dust (defined in the usual way such that the dust emissivity at frequency ν is $\propto \nu^\beta$), from ~ 1.5 in 2001 August to ~ 0 in 2003. This may indicate changes in (a) the nature of the dust (e.g. as grains grow, or are processed) and/or (b) the dust distribution, allowing hotter grains (which have flatter emissivities, cf. Mennella et al. 1998) to contribute more to the observed flux. However any conclusion along these lines must await detailed modelling combining both UKIRT (e.g. Tyne et al. 2002, Eyres et al., in preparation) and JCMT datasets; such an analysis will be reported elsewhere.


The referee has pointed out that the sub-millimetre rise could be due to the anticipated rise in effective temperature as the star retraces its post-AGB evolution towards the PN phase, and the consequent heating of the inner dust shell. While observational evidence that this process may have started has been presented by Kerber et al. (2002), we do not consider that this accounts for the rise in the sub-millimetre flux for reasons which may be discerned from Fig. 2.

In this Figure we plot the JCMT data from 15 June 2003 (Table 1), together with the 2–5 μ m data obtained in September 2003 from our UKIRT programme (Eyres et al., in preparation); we assume that variations are sufficiently slow that the data can be combined. Both sets of data are reasonably fitted by a $\sim 360 \pm 10$ K black-body (cf. Equation (2) below, and 350 ± 30 K for the UKIRT data alone; Eyres et al. 2004), suggesting that the 2–5 μ m and sub-millimetre emission have common origin.

We might envisage a scenario in which the sub-millimetre emission in Fig. 2 comes from inner, hotter, dust whose near-IR emission does not penetrate the outer region

Table 2. Properties of the dust in the environment of Sakurai's Object. A gas-to-dust ratio of 75 is assumed; see text for details.

Date D/M/Y	MJD −2450000	$M_d T_d$ ($10^{-5} M_\odot$ K)		T_d (K)	M_d ($10^{-7} M_\odot$)		M_{cs} ($10^{-5} M_\odot$)	
		450 μ m	850 μ m		450 μ m	850 μ m	450 μ m	850 μ m
18/08/2001	2140	4.6 ± 1.8	3.3 ± 1.1	504	0.9 ± 0.4	0.7 ± 0.2	0.7	0.5
09/01/2003	2649	2.7 ± 0.7	11.1 ± 0.8	418	0.6 ± 0.2	2.7 ± 0.3	0.5	2.0
05/05/2003	2765	7.5 ± 1.2	16.2 ± 0.8	401	1.9 ± 0.4	4.0 ± 0.5	1.4	3.0
15/06/2003	2806	10.2 ± 1.4	19.1 ± 0.9	395	2.6 ± 0.4	4.8 ± 0.5	1.9	3.6
16/06/2003	2807	—	18.2 ± 1.0	395	—	4.6 ± 0.5	—	3.5


Figure 2. Combined UKIRT/JCMT data for Sakurai's Object in late 2003 to give the 2.5–850 μ m spectral energy distribution; the UKIRT data have been binned for clarity. The curve is a 360 K black-body, fitted by eye. See text for details.


of the dust shell, while the observed near-IR emission comes from dust in the outer shell having very steep β -index. However such a situation would be *ad hoc* and contrived. Indeed that the β -index is close to zero in late 2003 is supported both by the 450/850 flux ratio, and by the closeness of the 2–850 μ m data to a black-body at ~ 360 K (see Fig. 2).

We therefore make the assumptions that (i) the emission we detect at 450 μ m and 850 μ m is from the dust shell (any emission at these wavelengths from the star – whatever its effective temperature – is at the few μ Jy level at most), (ii) the dust shell is optically thin at sub-millimetre wavelengths and (iii) the sub-millimetre emission is on the Rayleigh-Jeans tail.

We justify (ii) by noting that the optical depth τ at wavelength λ (in μ m) is $\sim 0.55 \tau(0.55 \mu\text{m})/\lambda(\mu\text{m}) \ll 1$, since the optical depth in the visual is ~ 10 (see Tyne et al. 2002), and (iii) is valid if the dust temperature is $\gtrsim 200$ –300 K. Eyres et al. (2004) find that a 1–5 μ m spectrum of Sakurai's Object obtained on 2003 September 8 is well-described by a black-body at temperature $\sim 350 \pm 30$ K, consistent with continued cooling of the dust shell; thus (iii) is certainly justified. We therefore have that

$$M_d T_d = \frac{f_\nu D^2 \lambda^2}{2\gamma \kappa k} \approx \frac{A}{\gamma} \left(\frac{f_\nu}{\text{mJy}} \right) \left(\frac{D}{\text{kpc}} \right)^2 M_\odot \text{ K}, \quad (1)$$

where the constant $A = 9.60 \times 10^{-7}$ for 450 μ m, and 6.23×10^{-6} for 850 μ m. In Equation (1) M_d is the dust mass, D is the distance of Sakurai's Object (see Kimeswenger 2002), T_d


Figure 3. Increase in circumstellar mass around Sakurai's Object as deduced from 450 μ m data (squares) and 850 μ m data (triangles), for a gas-to-dust ratio of 75. Lines are included to guide the eye. See text for details

is an appropriate value for the dust temperature (see below), κ is the absorption coefficient for the dust, and γ depends on physical conditions in the dust shell (e.g. $\gamma \equiv 1$ for an optically thin, isothermal shell).

Although Equation (1) applies to an optically thin, isothermal shell, it is also accurate, to within a factor $\simeq 2/(1 + (R_i/R_o)^{1/2}) \leq 2$ (where R_i, R_o are respectively the inner and outer radii of the dust shell), to the case of a dust shell with grain density law $n(r) \propto r^{-2}$ that is optically thick at short ($\sim$ optical/near-IR) wavelengths, optically thin in the sub-millimetre, and for which the dust temperature depends on distance r from the star as $T_d \propto r^{-1/2}$ (see Ivezić & Elitzur 1997). In this case T_d in Equation (1) is the temperature of the coolest dust; we take this to be the dust seen in the 1–5 μ m data, which still detect the dust ‘pseudophotosphere’ (Tyne et al. 2002, Eyres et al., in preparation). It is not unreasonable to assume that the dust shell around Sakurai's Object is geometrically thick, so that $R_o/R_i \gg 1$; for this case $\gamma = 2$ and we use this value here (see also below).

For the distance, we take $D = 2$ kpc (see Kimeswenger 2002), and we have taken κ from Mennella et al. (1998; their κ values for amorphous carbon at 295 K, which is likely to be close to that appropriate for Sakurai's Object).

We give values of $M_d T_d$, computed from Equation (1), in Table 2 for both wavelengths. For each epoch, we have estimated the temperature T_d of the coolest dust by (i) fitting a black-body to 1–5 μ m data obtained in the course of our IR monitoring programme (see Tyne et al. 2002, Eyres et al. 2004, and paper in preparation) and (ii) fitting an exponential decline to the time-dependence of T_d . We find:

$$T_d \text{ (K)} \simeq 817 \exp \left[\frac{(815.5 - t')}{2735.3} \right], \quad (2)$$

where $t' = \text{MJD} - 2450000$ and the uncertainty in T_d is typically $\lesssim 10\%$ in reproducing the values in Tyne et al. (2002) and Eyres et al. (2004). We stress that this expression is not physically meaningful, and we will refine it as we obtain further IR data; we use it simply as a means of interpolating the value of T_d at the time of our SCUBA observations. These values are included in Table 2, together with the corresponding dust masses M_d ; the uncertainties in M_d include an uncertainty of 10% in estimating T_d .

Neither the values of $M_d T_d$, nor of M_d , should of course depend on wavelength, and the discrepancies between the $450 \mu\text{m}$ and $850 \mu\text{m}$ values are due to the uncertainties in the $450 \mu\text{m}$ calibration, the error in the interpolated temperature, uncertainty in the properties of the dust, and the fact that the $1 - 5 \mu\text{m}$ emission differs somewhat from a black-body. Nevertheless the data at both wavelengths are consistent with a steady increase in the dust mass between 2001 August and 2003 June.

The dust mass may be converted to a circumstellar mass M_{cs} if we assume a gas-to-dust ratio, and that the gas-to-dust ratio remains constant in time. The gas-to-dust ratio for solar abundances and in the interstellar medium is $\simeq 120$ (e.g. Whittet 2003), while a value 200 was assumed by Tyne et al. (2002). These values are unlikely to be appropriate for the H-deficient, He- and C-rich wind of Sakurai's Object. If we take the abundances for 1996 July 3 from Asplund et al. (1999), assume that all O is locked up in CO (see Eyres et al. 2004) and that the remaining C is tied up in dust, we get a gas-to-dust ratio of 26; this is most likely a lower limit.

We assume an intermediate value of 75 here; the uncertainties in the gas-to-dust ratio and in the value of γ clearly lead to a corresponding uncertainty in dust mass, but we believe that *changes* in the latter should be reliable. The corresponding circumstellar masses are given in Table 2; the dependence of circumstellar mass on time is shown in Fig. 3.

We use these data to determine the average mass-loss rate over the period 2001 August to 2003 June. Using the $850 \mu\text{m}$ data, we find that $\dot{M} \simeq 1.6 \pm 0.3 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ over this period; the corresponding value from the $450 \mu\text{m}$ data is $4.9 \pm 4.4 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$. Taking the 2003 observations only, the $450 \mu\text{m}$ and $850 \mu\text{m}$ data give $\dot{M} \simeq 3.3 \pm 0.3 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ and $\dot{M} \simeq 3.6 \pm 0.3 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ respectively, with a weighted mean of $\dot{M} \simeq 3.4 \pm 0.2 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ (cf. Fig. 3); for comparison, the maximum mass-loss rate for post-AGB stars is $\sim 1 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$ (van Winckel 2003).

Tyne et al. (2002) used the DUSTY code (Ivezić & Elitzur 1997) to fit the $1 - 5 \mu\text{m}$ data with a carbon dust shell and, adjusted for a gas-to-dust ratio of 75, found that $\dot{M}$ increased from $1.8 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$ to $4.4 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$ over the period 1999 May to 2001 September. Our results suggest that the mass-loss from Sakurai's Object continues to rise.

4 CONCLUSIONS

We have presented $450 \mu\text{m}$ and $850 \mu\text{m}$ photometry of Sakurai's Object, and find that the flux densities at both wave-

lengths have increased steadily over the period 2001 August to 2003 June. These data, combined with dust temperatures obtained from our IR monitoring programme, lead to a mass-loss in 2003 of $\sim 3.4 \pm 0.2 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$, consistent with a continuing rise since our IR observations of 1999-2003.

Monitoring of this object at sub-millimetre wavelengths is continuing and further observations will be presented elsewhere.

ACKNOWLEDGMENTS

We thank the referee, Stefan Kimeswenger, for comments that helped to clarify some of the discussion. TRG is supported by the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., on behalf of the international Gemini partnership of Argentina, Australia, Brazil, Canada, Chile, the United Kingdom, and the United States of America. The JCMT is operated by the Joint Astronomy Centre on behalf of the Particle Physics and Astronomy Research Council (PPARC) of the United Kingdom, the Netherlands Organization for Scientific Research and the National Research Council of Canada. We thank the PPARC Panel for the Allocation of Telescope Time for supporting of this project. Data reduction was carried out using hardware and software provided by PPARC.

REFERENCES

- Asplund M., Lambert D. L., Kipper T., Pollacco D., Shetrone M. D., 1999, *A&A*, 343, 507
- Bond H. E., Pollacco D., 2002, *Keele Workshop on Sakurai's Object*, Eds A. Evans & B. Smalley, *Ap&SpSci*, 279, 31
- Evans A., Geballe T. R., Tyne V. H., Pollacco D., Eyres S. P. S., Smalley B., 2002, *MNRAS*, 332, L69 (Paper I)
- Eyres S. P. S., Smalley B., Geballe T. R., Evans A., Asplund M., Tyne V. H., 1999, *MNRAS*, 307, L11
- Eyres S. P. S., Geballe T. R., Tyne V. H., Evans A., Smalley B., Worters H. L., 2004, *MNRAS*, 350, L9
- Geballe T. R., Evans A., Smalley B., Tyne V. H., Eyres S. P. S., 2002, *Keele Workshop on Sakurai's Object*, Eds A. Evans & B. Smalley, *Ap&SpSci*, 279, 39
- Herwig F., 2001, *ApJ*, 554, L71
- Holland W. S., et al., 1999, *MNRAS*, 303, 659
- Ivezić Z., Elitzur M., 1997, *MNRAS*, 287, 799
- Jenness T., Lightfoot J. F., 2000, *Starlink User Note* 216, Starlink Project, CLRC
- Kerber F., Pirzcal N., de Marco O., Asplund M., Clayton G. C., Rosa M. R., 2002, *ApJ*, 581, L39
- Kimeswenger S., 2002, *Keele Workshop on Sakurai's Object*, Eds A. Evans & B. Smalley, *Ap&SpSci*, 279, 79
- Mennella V., Brucato J. R., Colangeli L., Palumbo P., Rotundi A., Bussoletto E., 1998, *ApJ*, 496, 1058
- Pollacco D., 1999, *MNRAS*, 304, 127
- Tyne V. H., Evans A., Geballe T. R., Eyres S. P. S., Smalley B., Dürbeck H. W., 2002, 334, 875
- van Winckel H., 2003, *ARAA*, 41, 391
- Whittet D. C. B., 2003, *Dust in the Galactic environment*, 2nd edition, Institute of Physics Publishing, Bristol & Philadelphia

This paper has been typeset from a $\text{\TeX}$ / $\text{\LaTeX}$ file prepared by the author.